1137

REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

wing instructions, searching existing data sources, ing this burden estimate or any other aspect or the management (215 series as America, 1215 series as America, 1215 series as 1215 series 1215 se

Dove Highway, Suite 1201, Arthquen, VA 22202-4302	, and to the Office of Management s		
1. AGENCY USE ONLY (Leave blank)	Jan 1983	3. REPORT TYPE AN Final (1 De	D DATES COVERED ac 81- 30 Nov 82
4. TITLE AND SUSTITLE			5. FUNDING NUMBERS
THREE-DIMENSIONAL INTER BOUNDARY LAYER FLOWS"	N HIGH SPEED	61102F	
6. AUTHOR(S)			2307/K1
G.R. Inger			
7. PERFORMING ORGANIZATION NAME University of Colorado	•		B. PERFORMING ORGANIZATION REPORT NUMBER
Dept of Aerospace Engine Boulder, CO 80309	eering	् अल ङ्काः ।	50 04 1 501
9. SPONSORING/MONITORING AGENC	NAME(S) AND ADDRESS	ES)	10. SPONSORING / MONITORING AGENCY REPORT NUMBER
AFOSR BLDG 410			
BAFB DC 20332-6448			AFOSR-82-0056
11. SUPPLEMENTARY NOTES	······································		*
12a. DISTRIBUTION / AVAILABILITY STA	TEMENT		12b. DISTRIBUTION CODE
Aphrove, for a distrall. 1999			
13. ABSTRACT (Maximum 200 words)			



14.	SUBJECT TERMS			15. NUMBER OF PAGES	
				16. PRICE CODE	
17	SECURITY CLASSIFICATION OF REPORT	18. SECURITY CLASSIFICATION OF THIS PAGE	19. SECURITY CLASSIFICATION OF ABSTRACT	20. LIMITATION OF ABSTRACT	
	unclassified	unclassified	1	<u> </u>	

NSN 7540-01-280-5500

12 01 112 89

Standard Form 298 (890104 Draft) Protested by ANN Ind. 239-18 298-01

Final Report on the First Year Effort

AFOSR 2-0056

" Three-Dimensional Interaction Effects in High Speed Boundary Layer Flows "

Dr. G.R. Inger
University of Colorado at Boulder

Jan. 3, 1983

ica.	Tro-grand	
Fire		
		- 1
i s		
	transis in the second second	ا ا
· ·	The second secon	-
A-1		1

1. Introduction

The objective of this research is the basic theoretical investigation of three-dimensional pressure, skin friction and heat transfer disturbances in both laminar and turbulent boundary layer flows including viscous-inviscid interaction effects, separation and reattachment. A sound understanding of these phenomena is required in modern aerodynamic design analyses of high-speed flight vehicles.

The aforementioned need has led to a systematic basic research effort treating in parallel the twin problems of (1) viscous-inviscid interaction with flow separation and/or attachment and (2) three-dimensional effects (including those due to streamwise vortices) in high-speed laminar and turbulent boundary layer flows. Such a study was initiated October 1, 1971 and has continued under AFOSR support. To date, it has been very successful both in solving a number of the target flow problems and in developing new ideas and tools 1-5. In particular, a premising three-dimensional extension of successful non-asymptotic triple deck theory for two-dimensional shock - turbulent boundary layer interactions was developed in collaboration with the Princeton Group's experimental studies 7,8 of the 3-B interaction phenomena illustrated in Fig. 1. This report summarines our recent progress during the past year.

2. Summary of First Year's Progress

The project initiated in 1981 was understood to be a long range multi-year effort; in the initial year at the University of Colorado, we therefore concentrated on first strengthening and extending certain key aspects of our analytical triple-deck model approach in two-dimensional flow to expedite its subsequent application to the more general three-dimensional case. These activities have gone very well.

As part of a substantial involvement in the detailed research, the principal investigator familiarized himself with the Fast Fourier Transform ("FFT") technique and its operational mathematical content, since this may well be an important component in the actual solution techniques of the 3-D problem. Moreover, we have succeeded in extending Lighthill's classical interaction theory (as well as Inger's generalization to turbulent flow) to include wall suction or blowing effects; the results, which are in the process of final completion, provide for the first time a really fundamentallybased analytic theory for understanding and calculating these important effects in a variety of specific supersonic SELI problems in either external aerodynamic or internal inlet flows. Comparisons will be made with available experimental studies such as those carried out at Cambridge, NASA-Lewis and (especially) at the University of Poitiers by LeBlanc, with whom the author has established collaboration under the aegis of an earlier NATO travel grant. We intend to prepare several major technical papers on this work for future technical testing presentation and journal publication.

We have been fortunate to also find an approximate but accurate closedform solution to the triple-deck equations including the turbulent case
which describes all the major physical features of the interaction zone (see
Fig. 2); this was achieved by implementing a small (but NOT negligible) lateral
pressure gradient approximation of the yields a very elegant and revealing analytical solution. Even better, this approximation appears adatable to the 3-D case
we well, which would considerably simplify its more formidable numerical aspects while illuminating the 3-D interactive physics involved. Furthermore,
we carried out a detailed study of the available turbulence eddy-viscosity
models suitable to the Law of the Wall region in three-dimensional turbulent

boundary layers (this is very important to the inner deck part of our 3-D triple deck approach). One consequence of this was the formulation of a new 3-D Law of the Wall theory for <u>incipiently-separating</u> conditions, which is the appropriate generalization of the Stratford wall velocity profile model at incipient separation in 2-D flow. A parametric study of the resulting universal non-dimensional profile solutions as a function of the basic spanwise-streamwise pressure gradient ratio parameter involved was completed.


During this activity, we maintained a good technical collaboration with Dr. Settles of Princeton including a day-long visit to his laboratory in April. We are making our theoretical developments available to assist, for example, interpretation of his 3-D upstream influence measurements on the swept ramp problem while he in turn has kept us up to date on the experimental findings.


As part of this research effort, technical liaison has also been maintained with certain relevant investigators in Western Europe with whom mutual scientific enchange has proven fruitful in the past (e.g., Professor Klaus Gersten of the Ruhr Universität Bochum-FRG and Dr. H. Stanewsky of the DFVLR-AVA in Göttingen-FRG, Drs. LeBlanc and Ardonceau of the CEAT in Poitiers, France, and Professor Keith Stewartson and Sir M. J. Lighthill of the University of London. Valuable technical discussion of viscous-inviscid interaction phanomena were held with these people during a trip to Europe last March. Moreover, a good contact was established with Dr. Ray Stalker (University of Sydney) who is a pioneer worker on the 3-D interaction problem.

References

- 1. Inger, G. R. "Three-Dimensional Disturbances in Reattaching Separated Flow," Proc. AGARD Symposium on Flow Separation, CP-168, pp. 18-1 to 18-12.
- 2. Inger, G. R., "Three-Dimensional Heat and Mass Transfer Effects Across High Speed Reattaching Flows," VPI&SU Report Aero-036, Blacksburg, Virginia, July 1975. Presented as AIAA Paper 76-160, 14th Aerospace Sciences Meeting, Washington, D.C., January 1976; AIAA Journal 15, March 1977, pp. 383-89.
- 3. Namtu, M., and Inger, G. R., "Spanwise-Periodic Three-Dimensional Disturbances in Nominally 2-D Separating Laminar Boundary Layer Flows," VPI&SU Report Aero-052, Blacksburg, Virginia, August 1976.
- 4. Inger, G. R., "On the Curvature of Compressible Boundary Layer Flows Near Separation," Zeitschrift für Agenwante Mekanik und Physik ("ZAMP"), Vol. 28/6, 1977, pp. 1027-1035.
- 5. Inger, G. R., "Compressible Laminar and Turbulent Boundary Layers Near Separation or Reattachment," <u>AIAA Paper 78-1164</u>, 11th Fluid and Plasma Dynamics Conference, Seattle, July 1978.
- 6. Inger, G. R., "The Skin Friction-Disturbance Sublayer in Non-Separating Shock-Turbulent Boundary Layer Interactions, AIAA Paper 80-1411, 13th AIAA Fluid and Plasma Dynamics Conference, Snowmass Colorado, July 14-16, 1980.
- 7. Oskam, B., "Three-Dimensional Flowfields Generated by the Interaction of a Swept Shock Wave With a Turbulant Brundary Layer," Ph. D. Thesis, MAE Department, Princeson University, 1976.
- 8. Settles, G. S., and Perkins, J. J., "Investigation of Three-Dimensional Shock/ Boundary Layer Interactions at Swept Compression Corners," AIAA Paper 79-1498.
- 9. Lighthill. M. J., "On Boundary Layers and Upstream Influence. II Supersonic Flows Without Separation," <u>Proc. Royal Soc. A 217</u>, 1953, pp. 478-507.
- 18. Stratford. B. B. "The Prediction of Separation of the Turbulent Boundary Layer,"


 Jour. Fluid Nech. 5, 1959, pp. 1-16.
- 11. Grew. S. C.. The Spanwise Perturbation of Two-Dimensional Boundary Layers," Jour. Fluid Nech. 24. Pt. 1, 1966, pp. 153-164.
- 12. Fannalop, T. M.. "Effects of Streamwise Vortices on Laminar Boundary Layer Flow," Trans. ASME Jour. App. Mech., June 1968, pp. 424-426.





B. SWEPT RAMP

FIG. 1



Triple-Deck Structure of the Interaction (Schematic)